

90-092670

B3



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

0 360 018  
A1



## EUROPEAN PATENT APPLICATION

Application number: 89115516.0

(91) Int. Cl.<sup>5</sup>: C07H 19/06, A61K 31/70

(12) Date of filing: 23.08.89

(20) Priority: 25.08.88 JP 212061/88

(43) Date of publication of application:  
28.03.90 Bulletin 90/13(54) Designated Contracting States:  
AT BE CH DE ES FR GB IT LI NL SE(71) Applicant: YOSHITOMI PHARMACEUTICAL  
INDUSTRIES, LTD.  
6-9, Hiranomachi 2-chome Chuo-ku  
Osaka-shi Osaka 105(JP)Applicant: YAMASA SHOYU CO., LTD.  
10-1, Araoi-Cho 2-Chome  
Choshi-Shi Chiba-Ken 288(JP)(72) Inventor: Ueda, Tohru  
6-27, Maruyamanishimachi 8-chome Chuo-ku  
Sapporo-shi Hokkaido 064(JP)  
Inventor: Sasaki, TakumaHelwashukusha (C)-57-11 18-15, Helwacho  
3-chome

Kanazawa-shi Ishikawa 921(JP)

Inventor: Matsuda, Akira

Minamishinkawa-Komuinshukusha

10-501, Nishi 13-chome Kita 23-jo Kita-ku

Sapporo-shi Hokkaido 001(JP)

Inventor: Yamagami, Keiji

407-14-103, Oaza-Kamifujisawa

Iruma-shi Saitama 358(JP)

Inventor: Fujii, Akhiro

12-19, Kamikkyoto 2-chome

Kiyose-shi Tokyo 204(JP)

(74) Representative: von Kreisler, Alek et al  
Patentanwälte Von  
Kreisler-Selting-Werner-Schönwald-Fues-Da-  
lmeyer Delchmannhaus  
D-5000 Köln 1(DE)

(52) 2'-Methylidenepyrimidin nucleoside compounds, their use and method for production thereof.


(57) 2'-Methylidenepyrimidine nucleoside compounds of the general formula:

NEW 2'-METHYLIDENE-2'-DEOXY-

CYTIDINE AND URIDINE

DERIVS. + USE AS

ANTITUMOUR AGENTS



90092670

Xerox Copy Centre

EP 0 360 018 A1

453

B

BEST AVAILABLE COPY

wherein R<sup>1</sup> stands for amino or hydroxy group; R<sup>2</sup> stands for a halogen or a lower alkyl when R<sup>1</sup> is amino or R<sup>2</sup> stands for an alkyl having 2 to 4 carbon atoms, an alkynyl having 2 to 4 carbon atoms or a haloalkyl when R<sup>1</sup> is hydroxy group; and R<sup>3</sup> stands for hydrogen or a phosphoric acid residue, or salts thereof, anticancer compositions containing one or more of these compounds and methods for production of these compounds.

Said compounds and salts thereof exhibit noticeable antitumor activities and are useful as anticancers.

90092670

## 2'-Methylidenepyrimidine Nucleoside Compounds, Their Use and Method for Production Thereof

## BACKGROUND OF THE INVENTION

5 This invention relates to novel 2'-methylidenepyrimidine nucleosides possessing an excellent antitumor action, their pharmaceutical use and a method for producing said compounds.

Under the circumstances in which death due to cancer has increased in number, chemotherapy and immunotherapy in addition to surgical therapy have been widely conducted. In this connection, in chemotherapy, cytosine arabinoside, 5-fluorouracil and the like as antimetabolites, which are considered  
 10 effective against acute leukemia, have been clinically used. However, the hitherto-known anticancer agents leave much to be desired in respect to therapeutic effect and involve various problems such as side effects. Thus, the development of excellent anticancer agents has been desired from various fields. Under such situations, the present inventors created 2'-deoxy-2'-methylidenecytidine and its pharmaceutically acceptable salts. However, since 2'-deoxy-2'-methylidenecytidine is inferior in solubility and its acid addition salts  
 15 are unstable, they are unsatisfactory in the aspect of pharmaceutical production.

## SUMMARY OF THE INVENTION

20 The object of the present invention is to provide novel compounds possessing excellent antitumor actions and their pharmaceutical use.

Another object of the present invention is to provide methods for producing said novel compounds.

The present inventors had conducted extensive studies to develop novel compounds useful as an  
 25 anticancer agent and having no problems in the aspect of pharmaceutical production and found that 2'-methylidenepyrimidine nucleoside compounds possessed excellent antitumor actions and no problems in the aspect of pharmaceutical production, which culminated in the completion of the present invention.


This invention relates to a 2'-deoxy-2'-methylidenepyrimidine nucleoside compound of the general formula :

30

35

40

45



(I)

50 wherein R<sup>1</sup> stands for amino or hydroxy group; R<sup>2</sup> stands for a halogen or a lower alkyl when R<sup>1</sup> is amino or R<sup>2</sup> stands for an alkyl having 2 to 4 carbon atoms, an alkynyl having 2 to 4 carbon atoms or a haloalkyl when R<sup>1</sup> is hydroxy group; and R<sup>3</sup> stands for hydrogen or a phosphoric acid residue, or a salt thereof.

Further, the present invention relates to an anticancer composition characterized by containing, as the active ingredient, a compound of the general formula (I) or a salt thereof.


Furthermore, the present invention relates to a method for producing compound (I).

In the above-mentioned general formula (I), with reference to  $R^2$ , the halogen means fluorine, chlorine, bromine or iodine; the lower alkyl means methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl or the like; the alkyl having 2 to 4 carbon atoms means ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl or the like; the alkynyl having 2 to 4 carbon atoms means ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl or the like; and as the halogen of the haloalkyl, mention is made of the halogens as mentioned above and as the alkyl of the haloalkyl, mention is made of the lower alkyls as mentioned above, and such haloalkyls include, for example, trifluoromethyl and chloroethyl.

As the salts of the compounds of the general formula (I), in case where  $R^3$  is hydrogen, mention is made of acid addition salts such as hydrochlorides, sulfates, hydrobromides, phosphates, maleates, fumarates, tartrates, succinates, citrates and p-toluenesulfonates, and in case where  $R^3$  is a phosphoric acid residue, mention is made of pharmaceutically acceptable salts exemplified by metal salts such as sodium salt, potassium salt, lithium salt, and calcium salt, or ammonium salt.

The present invention encompasses hydrates (1 hydrate, 1/2 hydrate, 1/4 hydrate, 3/2 hydrate) or other solvates of the compounds of the general formula (I) or their salts.

In the case of the compounds of the general formula (I) wherein  $R^1$  is hydroxy group, the present invention encompasses the uridine compounds of the general formula :




wherein  $R^2$  and  $R^3$  are of the same meaning as mentioned above, which are the tautomers of the compounds of the general formula (I) wherein  $R^1$  is hydroxy group.

As the compound of the general formula (I), there can be mentioned, for example, 2'-deoxy-2'-methylidene-5-fluorocytidine, 2'-deoxy-2'-methylidene-5-chlorocytidine, 2'-deoxy-2'-methylidene-5-bromocytidine, 2'-deoxy-2'-methylidene-5-iodocytidine, 2'-deoxy-2'-methylidene-5-methylcytidine, 2'-deoxy-2'-methylidene-5-ethylcytidine, 2'-deoxy-2'-methylidene-5-ethyluridine, 2'-deoxy-2'-methylidene-5-ethynyluridine or 2'-deoxy-2'-methylidene-5-fluorocytidine-5-phosphoric acid.

The compounds of the general formula (I) of the present invention wherein  $R^1$  is hydroxy group (hereinafter represented as the uridine compounds) can be produced by, for example, the following method:

After the hydroxy groups at the 3'- and 5'-positions of the sugar moiety of a compound of the general formula :

90092670



wherein R represents a halogen, an alkyl, an alkynyl having 2 to 4 carbon atoms or a haloalkyl are protected, the hydroxy group at the 2'-position of the sugar moiety is oxidized, followed by reaction with a Wittig reagent to obtain the compound having methyldene at the 2'-position of the sugar moiety and thereafter the protective groups of the hydroxy groups at the 3'- and 5'-positions of the obtained compound are removed, whereafter, if desired, the 5'-position of the sugar moiety is subjected to phosphorylation.

In the above-mentioned reactions, as the protective groups for the hydroxy groups at the 3'- and 5'-positions, any protective groups that are conventionally used as the protective groups for hydroxy groups can be applied. As such protective groups, for example, mention can be made of acyl groups such as acetyl, propionyl, butyryl, benzoyl and naphthoyl; acetal- or ketal-type protective groups such as ethyldene, propyldene, isopropyldene, benzylidene, cyclohexyldene, cyclopentyldene, methoxymethyldene, ethoxymethyldene and dimethoxyethyldene; aralkyl groups such as benzyl, p-methoxybenzyl, 3,4-dimethoxybenzyl, diphenylmethyl, triphenylmethyl,  $\alpha$ - or  $\beta$ -naphthylmethyl and  $\alpha$ -naphthylidiphenylmethyl; and silyl groups such as trimethylsilyl, t-butyldimethylsilyl, methyl-diisopropylsilyl, triisopropylsilyl and tetraisopropyl-disiloxy (TIPDS).

As the oxidation method for the hydroxy group at the 2'-position, there can be adopted chromic acid oxidation (A method) which comprises using a complex of chromic acid-pyridine-acetic anhydride and the like or activated dimethylsulfoxide oxidation (B method) which comprises using activated dimethylsulfoxide produced from oxalyl chloridedimethylsulfoxide. The oxidation reaction can be carried out in the presence of 1-10 times mole of an oxidizing agent relative to 1 mole of the compound to be oxidized at a temperature ranging from  $-10^{\circ}\text{C}$  to room temperature in the case of A method or at a temperature ranging from  $-10^{\circ}\text{C}$  to  $-80^{\circ}\text{C}$  in the case of B method for 1-10 hours.

As the Wittig reagent to be used for the methyldene reaction, specifically, triphenylphosphinemethylene can be mentioned, and preferably used is a compound which is prepared, immediately before use, from a triphenylphosphonium compound (methyltriphenylphosphonium bromide, methyltriphenylphosphonium iodide, etc.) and a strong alkali (potassium hydride, sodium hydride, n-butyl lithium, sodium amide, etc.) in accordance with the conventional methods. The Wittig reagent can be used suitably in an amount ranging from 1 to 3 times mole relative to 1 mole of the compound. The methyldene reaction with the use of a Wittig reagent can be carried out by reacting in a solvent (tetrahydrofuran, dioxane, ether, benzene, dimethylsulfoxide, etc., singly or in mixture) at  $-30$  to  $30^{\circ}\text{C}$  for 0.5 to 20 hours.

The deprotection reaction for the hydroxy groups of the sugar moiety can be conducted by suitably selecting conventional treatment such as acidic hydrolysis, alkaline hydrolysis, ammonium fluoride treatment or catalytic reduction in accordance with the protective group used. For example, when a silyl group is used as the protective group for the hydroxy group, the silyl group can be removed by ammonium fluoride treatment or acidic or alkaline hydrolysis.


The compounds wherein  $\text{R}^3$  is a phosphoric acid residue can be obtained by reacting with a phosphorylating agent (phosphorus oxychloride, tetrachloropyrophosphoric acid, etc.) which is conventionally used for selective phosphorylation at the 5'-position of nucleosides.

Moreover, the compounds of the present invention of the general formula (I) wherein  $\text{R}^1$  is amino can be prepared by subjecting a compound to an ordinary amination reaction converting a hydroxy group at the 4'-position of the base into an amino group, for example, by reacting a compound having methyldene at the

2'-position of the sugar moiety wherein the hydroxy groups at 3'- and 5'-positions of the sugar moiety are protected, which can be obtained by the above-mentioned method with 1,2,4-triazole in the presence of phosphorus oxychloride and triethylamine to obtain the 4-(1,2,4-triazol-1-yl) compound and converting the obtained compound to the 4-amino compound by passing ammonia gas thereto, followed by the removal of the protective groups for the hydroxy groups and, if desired, phosphorylation.

The reaction with 1,2,4-triazole proceeds under the flow of inert gas such as argon gas in a solvent such as acetonitrile at a temperature ranging from about 0°C to about room temperature for about 30 minutes to about 5 hours. The ammonia gas is preferably in advance passed through sodium hydroxide. Preferably, the amination reaction with ammonia gas is usually conducted for 1-5 hours.

Among the compounds of the general formula (II) to be used as the starting compounds in the present invention, the compounds of the general formula (II) wherein R is an alkynyl having 2-4 carbon atoms can be produced by, for example, the following method.



wherein R' represents an alkynyl having 2 to 4 carbon atoms protected by a silyl group and R'' represents an alkynyl having 2-4 carbon atoms.

In the above-mentioned reaction, a compound of the general formula (III) is reacted with a metal compound of silyl-protected alkynyl to obtain the compound (IV), followed by the removal reaction of the silyl group to obtain the objective compound (II').

Among the compounds of the general formula (II), the compounds wherein R is an alkyl having 2-4 carbon atoms can be produced by protecting the hydroxy groups at the 3'- and 5'-positions of the sugar moiety of the general formula (II) in accordance with the above-mentioned method and subjecting the protected compound to reduction reaction of the protected compound, followed by the removal of the protective groups of the hydroxy groups at the 3'- and 5'-positions of the sugar moiety.

When the compound of the present invention of the general formula (I) wherein R<sup>2</sup> is an alkyl having 2-4 carbon atoms is produced, without the removal of the hydroxy groups at the 3'- and 5'-positions of the sugar moiety of the above-mentioned reductive compounds, the compound having methyldene at the 2'-position of the sugar moiety is obtained in accordance with the methyldination hereinbefore mentioned, whereafter the protective groups are removed, followed by, if desired, phosphorylation at the 5'-position of the sugar moiety.

The objective compounds and synthetic intermediate compounds of the present invention can be isolated and purified by conventional purification and isolation methods (e.g. various chromatography method, ion exchange and adsorption chromatography, recrystallization, etc.) suitably in combination. Specifically, for example, in the case of the nucleoside compounds, after the solvent is distilled off, they are, if necessary, subjected to column chromatography and crystallized from a suitable organic solvent. In the case of nucleotide compounds (wherein R<sup>3</sup> is a phosphoric acid residue), they are purified by ion exchange chromatography with ion exchange resin, etc., adsorption column chromatography with activated charcoal, etc. or so on and lyophilization or crystallization to obtain the objective or intermediate compounds in the free acid form, or, if desired, as the salts.

The thus-obtained compounds of the general formula (I) can be converted into the above-mentioned acid addition salts, metal salts or ammonium salts.

Since the compounds of the present invention exhibit remarkable multiplication-suppressive actions

against tumor cells and have low toxicity, the pharmaceuticals of the present invention containing them as the active ingredient are useful as anticancer agents. The anticancer compositions of the present invention can be in forms such as powders, granules, tablets, sugar-coated tablets, capsules, syrups, suppositories, medicines for external use, injections and medicines for instillation in a mixture of an effective amount of the compound (I) and pharmaceutically acceptable carriers, excipients, diluents and the like.

While 2'-deoxy-2'-methylidenecytidine is not soluble in saline at the concentration of 25 mg/ml at room temperature, the compound of Example 1 is easily soluble under the same conditions.

The anticancer compositions of the present invention can be orally or non-orally administered to mammals including human. Any dosage is preferable so long as the compounds can inhibit the objective cancers effectively at the dosage. While the dosage varies depending on the animal to be treated, the kind of the cancer, the administration route, the composition form and the like, the daily dosage is generally 10 - 400 mg/kg body weight, preferably 50 - 200 mg/kg body weight, in the case of medicine for oral administration and 1 - 20 mg/kg body weight, preferably 1 - 5 mg/kg body weight, in the case of injections. The administration frequency can be suitably selected in the range from once to 4 times a day.

The anticancer compositions of the present invention can be used in combination with the other anticancer agents, immunostimulant or other acceptable medicaments.

#### PREFERRED EMBODIMENTS:

The invention will be hereinbelow described more specifically by way of examples, but should not be construed as limiting its scope to them.

#### Example 1

(1) 5-Fluorouridine, 2.62 g, was dissolved in 30 ml of pyridine, 3.3 ml of 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane was added at 0°C, and the mixture was stirred at 0°C for 2 hours and then at room temperature for 1.5 hours. By adding 5 ml of water, the solvent was distilled off under reduced pressure. After 100 ml of ethyl acetate and 100 ml of water were added to partition the solution, the organic layer was washed with water twice and then, dried over sodium sulphate and the solvent was distilled off under reduced pressure. The residue was packed into silica gel column (diameter, 2.4x23 cm) and eluted with 25% ethyl acetate/hexane to afford 3',5'-O-tetraisopropylidisiloxanyl-5-fluorouridine as a syrupy substance. Yield: 4.24 g, 83.9%; Mass (m/z): 504(M<sup>+</sup>), 461(M<sup>+</sup>-isopropyl)

(2) The compound obtained in (1) above, 5 g, was dissolved in 10 ml of methylene chloride and mixed with 4 equivalents of chromic acid complex (4 g of chromium oxide, 6.67 ml of pyridine, and 4 ml of acetic anhydride in 100 ml of methylene chloride). The mixture was stirred at room temperature for 1 hour and added dropwise to 500 ml of ethyl acetate and the solution was filtered through a silica gel filter (diameter, 6.0x1.5 cm). The filtrate was distilled off under reduced pressure and the residue was partitioned by 200 ml of ethyl acetate and 200 ml of water. The organic layer was washed with water, dried over sodium sulphate, and the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 2.4x28 cm), eluted with 20% ethyl acetate-hexane, and crystallized from hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-oxo-5-fluorouridine. Yield: 4.3 g, 86%; M.p. 183-186°C; Mass(m/z): 502(M<sup>+</sup>)

Elementary analysis: for C<sub>21</sub>H<sub>35</sub>N<sub>2</sub>O<sub>7</sub>FSi<sub>2</sub>:

Calculated:	C, 50.17; H, 7.02; N, 5.57
Found :	C, 50.29; H, 7.33; N, 5.58

(3) Methyltriphenylphosphonium bromide, 8.57 g, was suspended in 80 ml of tetrahydrofuran under argon gas stream, 15.1 ml of n-butyllithium (1.58 mol/L hexane solution) was added dropwise at 0°C, and the mixture was stirred at room temperature for 1 hour. A solution of 3.02 g of the compound obtained in (2) above in 25 ml of tetrahydrofuran was added dropwise under argon gas stream at 0°C. After 20 minutes, the temperature was restored to room temperature, at which stirring was continued for 3 hours. Neutralization was conducted with 1 N ammonium bromide and the solution was concentrated. The solution was partitioned with 100 ml of ethyl acetate and 100 ml of water, and the organic layer was washed with water.

dried over sodium sulphate and distilled off under diminished pressure. The residue was placed into a silica gel column (diameter, 3.5x18 cm) and eluted with 20% ethyl acetate/hexane to afford a syrupy product of 3,5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-fluorouridine. Yield: 2.43 g, 81%; Mass (mvz): 500(M<sup>+</sup>)

(4) Triethylamine, 1.4 ml, and 684 mg of 1,2,4-triazole were dissolved in 10 ml of acetonitrile and the solution was cooled under argon gas stream to 0 °C. Phosphorus oxychloride, 280 µl, was added dropwise, and the mixture was stirred at room temperature for 1 hour. The precipitate was filtered off, and to the filtrate was added dropwise a solution of 500 mg of the compound obtained in (3) above in 5 ml of acetonitrile. After 3 hours at room temperature, the solution was stirred at 50 °C for 1 hour and ammonia gas which had been passed over sodium hydroxide was passed through for 4 hours. The solution was partitioned with 100 ml of chloroform and 100 ml of water, and the aqueous layer was washed with chloroform twice. The collected chloroform layer was dried over sodium sulphate and distilled off under diminished pressure. The residue was placed on a column of silica gel (diameter, 3.5x14 cm) and eluted with 5% ethanol/chloroform to give 3,5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-fluorocytidine as a syrupy substance. Yield: 435 mg, 87%; Mass(mvz): 499(M<sup>+</sup>)

(5) The compound obtained in (4) above, 400 mg, was dissolved in 10 ml of tetrahydrofuran, 1.8 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. After neutralization with acetic acid, the solvent was distilled off under reduced pressure. The residue was packed into a silica gel column (diameter, 2.4x18 cm), eluted with ethanol/chloroform, and crystallized from ethanol/ethyl acetate to give 2'-deoxy-2'-methylidene-5-fluorocytidine. Yield: 177 mg, 88.2%; M.p. 185-187 °C; Mass(mvz): 257(M<sup>+</sup>)

Elementary analysis: for C<sub>10</sub>H<sub>12</sub>N<sub>3</sub>O<sub>4</sub>F:

Calculated:	C, 46.70; H, 4.70; N, 18.34
Found :	C, 46.06; H, 4.67; N, 16.00

#### Example 2

(1) A 1.4 ml quantity of triethylamine and 684 mg of 1,2,4-triazole were dissolved in 10 ml of acetonitrile and the solution was cooled under argon gas stream to 0 °C. Phosphorus oxychloride, 280 µl, was added by drops and the mixture was stirred at room temperature for 1 hour. The precipitate was filtered off, and to the filtrate was added dropwise a solution of 498 mg of 3,5'-O-tetraisopropylidisiloxanyl-2'-methylidenethymidine obtained from thymidine in a similar procedure to Example 1 (1) to (3). After 3 hours, ammonia gas which had been passed over sodium hydroxide was passed through. After 3 hours, 100 ml of chloroform and 100 ml of water were added for partitioning and the resulting aqueous layer was washed with chloroform twice. The collected chloroform layer was dried over sodium sulphate and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x12 cm) and eluted with 5% ethanol/chloroform to give 3,5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-methylcytidine as a syrupy substance. Yield: 315 mg, 63.3%; Mass(mvz): 495(M<sup>+</sup>)

(2) The compound obtained in (1) above, 300 mg, was dissolved in 10 ml of tetrahydrofuran and 1.4 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. After neutralization with acetic acid, the solvent was distilled under diminished pressure. The residue was placed into a silica gel column (diameter: 2.4x17 cm), eluted with ethanol/chloroform and crystallized from ethanol/ethyl acetate to give 2'-deoxy-2'-methylidene-5-methylcytidine. Yield: 137 mg, 89.0%; M.p. 198-204 °C (decomposition); Mass(mvz): 253(M<sup>+</sup>)

Elementary analysis: for C<sub>11</sub>H<sub>15</sub>N<sub>3</sub>O<sub>4</sub>:

Calculate:	C, 52.17; H, 5.97; N, 16.59
Found :	C, 52.02; H, 5.99; N, 16.37



## Example 3

(1) 5-Chlorouridine, 10 g, was dissolved in 100 ml of pyridine, and 12.44 ml of 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane was added at 0°C, and the mixture was stirred for 3 hours. After stirring at room temperature for 2 hours, 5 ml of ethanol was added and the solvent was distilled off under diminished pressure. The residue was partitioned with 200 ml of ethyl acetate and 200 ml of water, and the resulting organic layer was dried over sodium sulphate and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.0x25 cm) and eluted with 25% ethyl acetate/hexane to give a syrupy product of 3',5'-O-tetraisopropylidisiloxanyl-5-chlorouridine. Yield: 14.63 g, 78.2%; Mass(m/z): 478-(M<sup>+</sup>-isopropyl)

(2) The compound obtained in (1) above, 5.21 g, was dissolved in 15 ml of methylene chloride and mixed with 4 equivalents of chromic acid complex (4 g of chromium oxide, 6.6 ml of pyridine and 4 ml of acetic anhydride in 100 ml of methylene chloride), and the mixture was stirred for 2 hours. The reaction solution was added dropwise to 400 ml of ethyl acetate and filtered through a silica gel filter (dia meter, 6.0x1.0 cm). After the solvent was distilled off under reduced pressure, the residue was placed into a silica gel column (diameter, 2.4x30 cm), eluted with 25% ethyl acetate/hexane, and crystallized from ethyl acetate/hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-oxo-5-chlorouridine. Yield: 3.75 g, 72%; M.p. 139-201°C; Mass(m/z): 476(M<sup>+</sup>-isopropyl)

Elementary analysis: for C<sub>21</sub>H<sub>35</sub>N<sub>2</sub>O<sub>7</sub>ClSi<sub>2</sub>:

Calculated:	C, 48.49; H, 6.78; N, 5.39
Found :	C, 48.24; H, 7.02; N, 5.18

25

(3) Methyltriphenylphosphonium bromide, 5.72 g, was suspended in 50 ml of tetrahydrofuran under argon stream and 10.1 ml of n-butyllithium was added dropwise under ice cooling. At room temperature, the solution was stirred for 1 hour. Then, under cooling at 0°C, a solution of 2.08 g of the compound obtained in (2) above in 10 ml of tetrahydrofuran was added dropwise and after 30 minutes, the temperature was returned to room temperature. After 5 hours, neutralization was conducted with 1 N ammonium bromide and the solution was partitioned with 100 ml of ethyl acetate and 100 ml of water. The organic layer was washed with water, dried over sodium sulphate, and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 5.0x4 cm) and eluted with 25% ethyl acetate/hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-chlorouridine as a foamy product. Yield: 1.7 mg, 81.7%; Mass(m/z): 517(M<sup>+</sup>)

(4) A 3 ml quantity of triethylamine and 1.52 g of 1,2,4-triazole were dissolved in 20 ml of acetonitrile under argon stream and cooled with ice. To this was added dropwise 620 µl of phosphorus oxychloride and after 30 minutes, the solution was stirred at room temperature for 30 minutes. After the precipitate was filtered off, 1.04 g of the compound obtained in (3) above was added to the filtrate, and the mixture was stirred at room temperature for 3 hours. Ammonia gas which had been passed over sodium hydroxide was passed through the mixture for 2 hours, and partitioning with 100 ml of chloroform and 100 ml of water was conducted. The aqueous layer was washed with chloroform and the collected chloroform layer was dried over sodium sulphate and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x17 cm), eluted with 75% ethyl acetate-hexane and crystallized from ethyl acetate to give 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-chlorocytidine. Yield: 944 mg, 90.7%; M.p. 108-111°C; Mass(m/z): 517(M<sup>+</sup>)

(5) The compound obtained in (4) above, 900 mg, was dissolved in 10 ml of tetrahydrofuran, 4 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. After 5 ml of methanol was added, the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x12 cm), eluted with 20% ethanol/chloroform and crystallized from ethanol/ethyl acetate to give 2'-deoxy-2'-methylidene-5-chlorocytidine. Yield: 441 mg, 93.4%; M.p. 168-170°C; Mass(m/z): 273(M<sup>+</sup>)

55

## Example 4

461

90092670

(1) 5-Bromouridine, 3.32 g, was dissolved in 30 ml of pyridine and 3.3 ml of 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane was added at 0°C, and the mixture was stirred for 2 hours. After returning to room temperature, further stirring was conducted for 1 hour and 40 minutes. Water, 5 ml, was added to distill the solvent off under reduced pressure. After azeotropic distillation with ethanol, the residue was placed into a silica gel column (diameter, 2.4x25 cm) and eluted with 25% ethyl acetate/hexane to give 3',5'-O-tetraisopropylidisiloxanyl-5-bromouridine as a syrupy substance. Yield: 5.05 g, 89.2%; Mass(m/z): 522(M<sup>+</sup>-isopropyl)

(2) The compound obtained in (1) above, 3.57 g, was mixed with 4 equivalents of chromic acid complex (2.4 g of chromium oxide, 4 ml of pyridine and 2.4 ml acetic anhydride in 70 ml of methylene chloride) and stirred at room temperature for 40 minutes. The reaction solution was added dropwise to 400 ml of ethyl acetate and the solution was filtered through a silica gel filter (diameter, 6.0x1.0 cm). The filtrate was concentrated, and 150 ml of ethyl acetate and 150 ml of water were added for partitioning. The organic layer was washed with water and dried over sodium sulphate, and the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 5.0x14 cm), eluted with 20% ethyl acetate/hexane and crystallized from ethyl acetate-hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-oxo-5-bromouridine.

Yield: 3.01 g, 84.8%; M.p. 204-205°C; Mass(m/z): 520(M<sup>+</sup>-isopropyl)

Elementary analysis: for C<sub>21</sub>H<sub>35</sub>N<sub>2</sub>O<sub>7</sub>Si<sub>2</sub>Br:

Calculated:	C, 44.75; H, 6.26; N, 4.97
Found :	C, 44.57; H, 6.32; N, 4.92

(3) Methyltriphenylphosphonium bromide, 5.72 g, was suspended in 50 ml of tetrahydrofuran under argon stream, and 10.1 ml of n-butyllithium was added by drops at 0°C. The temperature was restored to room temperature. One hour thereafter, a solution of 2.25 g of the compound obtained in (2) above in 15 ml of tetrahydrofuran was added dropwise at 0°C. After 3 hours, the temperature was restored to room temperature and the solution was stirred overnight and neutralized with 1 N ammonium bromide. The solution was partitioned with 150 ml of ethyl acetate and 150 ml of water. The ethyl acetate layer was washed with water and dried over sodium sulphate, and the solvent was distilled off under diminished pressure. The residue was placed into a silica gel column (diameter, 5.0x14 cm) and eluted with 25% ethyl acetate/hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-bromouridine as a foamy substance. Yield: 1.72 g, 76.4%; Mass(m/z): 518(M<sup>+</sup>-isopropyl)

(4) Into 20 ml of acetonitrile were dissolved 3 ml of triethylamine and 1.52 g of 1,2,4-triazole under argon stream, and the solution was cooled with ice. To this was added dropwise 620 µl of phosphorus oxychloride, and after 30 minutes, the temperature was restored to room temperature. Thirty minutes thereafter, the precipitate was filtered off. To the filtrate was added 1.12 g of the compound of (3) above, and the mixture was stirred at room temperature for 3 hours. Ammonia gas which had been passed over sodium hydroxide was passed through the mixture for 3 hours, and 100 ml of chloroform and 100 ml of water were added to partition the solution. The water layer was washed with chloroform, the combined chloroform layer was dried over sodium sulfate, and the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x17 cm), eluted with 75% ethyl acetate-hexane and crystallized from ethyl acetate-hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-bromocytidine. Yield: 926 mg, 82.7%; M.p. 169-171°C; Mass(m/z): 560(M<sup>+</sup>)

(5) The compound of (4) above, 900 mg, was dissolved in 10 ml of tetrahydrofuran, 4 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. Methanol, 5 ml, was added, and the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x11 cm), eluted with 20% ethanol/chloroform, and crystallized from ethanol/ethyl acetate to obtain 2'-deoxy-2'-methylidene-5-bromocytidine. Yield, 518 mg, 102%; M.p. 172-180°C (decomposition); Mass(m/z): 318(M<sup>+</sup>)

#### Example 5

(1) 5-Iodouridine, 10 g, was dissolved in 100 ml of pyridine, 8.9 ml of 1,3-dichloro-1,1,3,3-tetraisopropyl-disiloxane was added at 0°C, and the mixture was stirred for 1.5 hours. The temperature was restored to

room temperature. Three hours thereafter, 10 ml of water was added and the solvent was distilled off under reduced pressure. After azeotropic distillation with ethanol, the residue was placed into a silica gel column (diameter, 3.0x30 cm) and eluted with 25% ethyl acetate/hexane to result in 3,5'-O-tetraisopropylidisiloxanyl-5-iodouridine as a syrupy substance. Yield, 14.7 g, 89%; Mass(m/z): 612(M<sup>+</sup>)

(2) The compound acquired in (1) above, 5.82 g, was mixed with 4 equivalents of chromic acid complex (3.55 g of chromium oxide, 5.9 ml of pyridine, and 3.55 ml of acetic anhydride in 100 ml of methylene chloride) and the mixture was stirred at room temperature for 40 minutes. The reaction solution was added by drops to 500 ml of ethyl acetate and filtered through silica gel (diameter, 6.0x15 cm), and the filtrate was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 5.0x21 cm) and eluted with 25% ethyl acetate/hexane to give 3,5'-O-tetraisopropylidisiloxanyl-2'-oxo-5-iodouridine. Yield, 1.72 g, 76.4%; Mass(m/z): 610(M<sup>+</sup>)

(3) Methyltriphenylphosphonium bromide, 7.94 g, was suspended in 80 ml of tetrahydrofuran under argon stream and 14 ml of n-butyllithium was added by drops at 0°C. The temperature was restored to room temperature. One hour thereafter, while the mixture was cooled to 0°C, a solution of 3.39 g of the compound obtained in (2) above in 15 ml of tetrahydrofuran was added dropwise. After one hour, the temperature was restored to room temperature, stirring was conducted overnight. After neutralization with 1 N ammonium bromide and concentration, the solution was partitioned with 200 ml of ethyl acetate and 200 ml of water. The organic layer was dried over sodium sulphate and distilled off under reduced pressure. The residue obtained was placed into a silica gel column (diameter, 5.0x22 cm) and eluted with 20% ethyl acetate/hexane to give 3,5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-iodouridine as a syrupy substance. Yield: 1.8 g, 47.2%; Mass(m/z): 608(M<sup>+</sup>)

(4) Triethylamine, 1.5 ml, and 760 mg of 1,2,4-triazole were dissolved in 10 ml of acetonitrile under argon gas stream and cooled with ice. Phosphorus oxychloride, 310  $\mu$ l, was added dropwise and 30 minutes thereafter, the temperature was restored to room temperature. After 30 minutes, the precipitate was filtered off and 321 mg of the compound of (3) above was added to the filtrate, followed by stirring at room temperature. Ammonia gas which had been passed over sodium hydroxide was introduced into the solution for 3 hours, and 50 ml of chloroform and 50 ml of water were added to partition the solution. The aqueous layer was washed with chloroform and the collected organic layer was dried over sodium sulphate and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 2.4x20 cm), eluted with 10% ethanol/chloroform and crystallized from ethyl acetate to give 3,5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-iodocytidine. Yield: 169 mg, 52.7%; M.p. 185-187°C; Mass(m/z): 607(M<sup>+</sup>)

(5) The compound obtained in (4) above, 160 mg, was dissolved in 5 ml of tetrahydrofuran, 1 ml of tetra-n-butyl-ammonium fluoride was added and the mixture was stirred at room temperature for 30 minutes. Methanol, 2 ml, was added and the solvent was distilled off under diminished pressure. The residue was placed into a silica gel column (diameter, 3.5x7.0 cm) and eluted with 20% ethanol/chloroform to give 2'-deoxy-2'-methylidene-5-iodocytidine as a syrupy product. Yield: 92 mg, 97.3%; M.p. 159-161°C; Mass(m/z): 365(M<sup>+</sup>)

#### Example 6

(1) 5-Ethynyluridine, 2.1 g, was dissolved in 50 ml of pyridine, 2.6 ml of 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane was added under ice cooling, and the mixture was stirred for 3 hours. At room temperature, stirring was conducted overnight, and 5 ml of methanol was added to distill the solvent off at reduced pressure. To the residue were added 100 ml of ethyl acetate and 100 ml of water for partitioning. The organic layer was washed with water twice and dried over sodium sulphate, and the solvent was distilled off under reduced pressure. The residue was placed into a column of silica gel (diameter, 5.0x17 cm) and eluted with 25% ethyl acetate/hexane to give 3,5'-O-tetraisopropylidisiloxanyl-5-ethynyluridine as a foamy product. Yield: 3.26 g, 82%; Mass(m/z): 510(M<sup>+</sup>)

(2) The compound obtained in (1) above, 2.04 g, was mixed with 4 equivalents of chromic acid complex (1.55 g of chromium oxide, 2.57 ml of pyridine, and 1.55 ml of acetic anhydride in 40 ml of methylene chloride) and the mixture was stirred at room temperature for 40 minutes. The reaction solution was added by drops to 200 ml of ethyl acetate and filtered through silica gel (diameter, 6.0x1.0 cm). The filtrate was distilled off at reduced pressure and the residue was partitioned with 100 ml of ethyl acetate and 100 ml of water. The organic layer was washed twice and dried over sodium sulphate, and the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x15 cm) and eluted

with 20% ethyl acetate/hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-oxo-5-ethynyluridine as a syrupy substance. Yield: 1.62 g, 81%; Mass(m/z): 508(M<sup>+</sup>)

(3) Methyltriphenylphosphonium bromide, 4.29 g, was suspended in 40 ml of tetrahydrofuran under argon gas stream, and under ice cooling, 7.58 ml of n-butyllithium was added dropwise. At room temperature, the solution was stirred for 1 hour and under ice cooling, a solution of 1.52 g of the compound obtained in (2) above in 10 ml of tetrahydrofuran was added dropwise. After 30 minutes, the temperature was restored to room temperature. Six hours thereafter, neutralization was conducted with 1 N ammonium bromide, and 100 ml of ethyl acetate and 100 ml of water were added to partition the solution. The organic layer was washed with water twice and dried over sodium sulphate, and then, the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x20 cm) and eluted with 20% ethyl acetate/hexane to result in 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-ethynyluridine as a syrupy substance. Yield: 1.12 g, 74%; Mass(m/z): 506(M<sup>+</sup>)

(4) The compound obtained in (3) above, 500 mg, was dissolved in 10 ml of tetrahydrofuran and 2.2 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 20 minutes. After neutralization with acetic acid, the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 2.4x18 cm), eluted with 10% ethanol/chloroform, and crystallized from ethanol-hexane to afford 2'-deoxy-2'-methylidene-5-ethynyluridine. Yield: 230 mg, 87%; M.p. 172-174°C; Mass(m/z): 264(M<sup>+</sup>)

20

## Example 7

(1) Triethylamine, 1.5 ml, and 760 mg of 1,2,4-triazole were dissolved in 10 ml of acetonitrile under argon gas stream and the solution was cooled with ice. A 310  $\mu$ l quantity of phosphorus oxychloride was added dropwise and 30 minutes thereafter, the solution was stirred at room temperature for 30 minutes. After the precipitate was filtered off, to the filtrate was added 500 mg of 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-ethynyluridine which was obtained in Example 8 (3), and the mixture was stirred at a room temperature for 3 hours. Ammonia gas which had been passed over sodium hydroxide was passed through for 2 hours, and 80 ml of chloroform and 80 ml of water were added for partitioning. The water layer was washed with chloroform and the collected chloroform layer was dried over sodium sulphate, followed by distillation off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x14 cm) and eluted with 75% ethyl acetate/hexane to give 3',5'-O-tetraisopropylidisiloxanyl-2'-deoxy-2'-methylidene-5-ethynylcytidine. Yield: 310 mg, 62%; Mass(m/z): 505(M<sup>+</sup>)

(2) The compound obtained in (1) above, 300 mg, was dissolved in 10 ml of tetrahydrofuran and 1.2 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. After 2 ml of methanol was added, the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 2.0x12 cm) and eluted with 20% ethanol/chloroform to result in 2'-deoxy-2'-methylidene-5-ethynylcytidine as a foamy product. Yield: 9.7 mg, 62%

40

## Example 8

(1) A 3.8 g quantity of 3',5'-O-tetraisopropylidisiloxanyl-5-ethynyluridine, which was obtained in Example 6 (1) was dissolved in 50 ml of methanol and 100 mg of 10% palladium carbon was suspended to conduct catalytic reduction under a moderate pressure of 20 p.s.i. for 4 hours. The reaction solution was filtered through cerite and the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.0x20 cm) and eluted with 33% ethyl acetate-hexane to give 3',5'-O-tetraisopropylidisiloxanyl-5-ethyluridine as a syrupy substance. Yield: 3.22 mg, 84%; Mass(m/z): 514(M<sup>+</sup>)

(2) The compound obtained in (1) above, 3.15 g, was mixed with 4 equivalents of chromic acid complex (2.37 g of chromium oxide, 3.93 ml of pyridine, and 2.37 ml of acetic anhydride in 60 ml of methylene chloride) and the mixture was stirred at room temperature for 50 minutes. The reaction solution was added dropwise to 300 ml of ethyl acetate and filtered through silica gel (diameter, 6.0x10 cm). After the filtrate was concentrated, 100 ml of ethyl acetate and 100 ml of water were added to partition the solution. The organic layer was washed with water twice and dried over sodium sulphate, followed by distillation off under reduced pressure. The residue was placed into a silica gel column (3.5 in diameter x 15 cm), eluted with 20% ethyl acetate-hexane and crystallized from ethyl acetate-hexane to give 3',5'-O-tetraisopropylidisiloxanyl-5-ethyluridine as a syrupy substance. Yield: 3.22 mg, 84%; Mass(m/z): 514(M<sup>+</sup>)

propyldisiloxanyl-2'-oxo-5-ethyluridine. Yield: 2.54 g, 80.8%; M.p. 176-177° C; Mass(m/z): 512(M<sup>+</sup>)  
 Elementary analysis; for C<sub>23</sub>H<sub>40</sub>N<sub>2</sub>O<sub>7</sub>Si<sub>2</sub>:

Calculated:	C, 53.88; H, 7.88; N, 5.48
Found :	C, 53.59; H, 8.01; N, 5.56

(3) Methyltriphenylphosphonium bromide, 5.72 g, was suspended in 50 ml of tetrahydrofuran under argon gas stream and under ice cooling. 10.1 ml of n-butyl-lithium was added dropwise. The mixture was stirred at room temperature for an hour. To the mixture was added dropwise the solution of 2.05 g of the compound obtained in (2) above in 15 ml of tetrahydrofuran under ice cooling. After 3 hours, the mixture was stirred at room temperature for 5 hours. The solution was neutralized with 1 N ammonium bromide, and 150 ml of ethyl acetate and 150 ml of water were added for partitioning. After washing with water twice, the organic layer was dried over sodium sulphate and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x13 cm), eluted with 20% ethyl acetate/hexane, and crystallized from ethyl acetate-hexane to give 3',5'-O-tetraisopropyldisiloxanyl-2'-deoxy-2'-methylidene-5-ethyluridine. Yield: 1.70 g, 82.7%; M.p. 153-155° C; Mass (m/z): 510(M<sup>+</sup>)  
 Elementary analysis: for C<sub>24</sub>H<sub>42</sub>N<sub>2</sub>O<sub>6</sub>Si<sub>2</sub>.

Calculated:	C, 56.44; H, 8.29; N, 5.48
Found :	C, 56.87; H, 8.41; N, 5.36

(4) The compound obtained in (3) above, 510 mg, was dissolved in 10 ml of tetrahydrofuran, and 2.2 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. After neutralization with acetic acid, the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.0x12 cm), eluted with 8.25% ethanol/chloroform, and crystallized from ethanol-hexane to give 2'-deoxy-2'-methylidene-5-ethyluridine. Yield: 234 mg, 87.2%; M.p. 137-139° C; Mass(m/z): 268(M<sup>+</sup>)  
 Elementary analysis; for C<sub>12</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>:

Calculated:	C, 53.73; H, 6.01; N, 10.44
Found :	C, 53.79; H, 5.89; N, 10.67

#### Example 9

(1) A 3 ml quantity of triethylamine and 1.52 g of 1,2,4-triazole were dissolved under argon stream in 20 ml of acetonitrile and cooled with ice. To this was added dropwise 620 μl of phosphorus oxychloride and after 30 minutes, the mixture was stirred at room temperature for 30 minutes. After the precipitate was filtered off, 1.2 g of 3',5'-O-tetraisopropyldisiloxanyl-2'-deoxy-2'-methylidene-5-ethyluridine which was obtained in Example 8 (3) was added to the filtrate and stirred at room temperature for 4 hours. Ammonium gas which was passed over sodium hydroxide was introduced into the solution and then, 100 ml of chloroform and 100 ml of water were added to partition the solution. The aqueous layer was washed with chloroform, and the combined chloroform was dried over sodium sulphate and distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x20 cm), eluted with 9% ethanol/chloroform, and crystallized from ethyl acetate/hexane to give 3',5'-O-tetraisopropyldisiloxanyl-2'-deoxy-2'-methylidene-5-ethylcytidine. Yield: 1.0 g, 83.3%; M.p. 197-200° C; Mass(m/z): 509(M<sup>+</sup>)

(2) The compound obtained in (1) above, 900 mg, was dissolved in 10 ml of tetrahydrofuran, 4 ml of tetra-n-butylammonium fluoride was added, and the mixture was stirred at room temperature for 30 minutes. After addition of 5 ml of methanol, the solvent was distilled off under reduced pressure. The residue was placed into a silica gel column (diameter, 3.5x10 cm), eluted with 20% ethanol/chloroform, and crystallized

from ethanol-ethyl acetate to give 2'-deoxy-2'-methylidene-5-ethylcytidine. Yield: 402 mg, 85.4%; M.p. 194-196° C; Mass(m/z): 267(M<sup>+</sup>)

## Example 10

2'-Deoxy-2'-methylidene-5-fluorocytidine and phosphorus oxychloride are made to react and treated in a usual manner to give 2'-deoxy-2'-methylidene-5-fluorocytidine-5'-phosphoric acid.

## Pharmaceutical Preparation Example 1

Tablets comprising:

Compound of Example 1	50.0 mg
Finely divided cellulose	25.0 mg
Lactose	49.5 mg
Starch	40.0 mg
Talc	5.0 mg
Magnesium stearate	0.5 mg

The tablets may be subjected to sugar coating or film coating treatment, if desired, thereby to obtain sugar-coated tablets or film-coated tablets.

## Pharmaceutical Preparation Example 2

Capsules encapsulated with the ingredients:

Compound of Example 1	50.0 mg
Lactose	50.0 mg
Starch	15.0 mg
Talc	5.0 mg

## Pharmaceutical Preparation Example 3

Fine powder comprising:

Compound of Example 1	10%
Lactose	80%
Starch	10%

## Pharmaceutical Preparation Example 4

Granules comprising:

90092670

Compound of Example 1	10%
Lactose	55%
Finely divided cellulose	20%
Starch	15%

10 Pharmaceutical Preparation Example 5

Compound of Example 1	50.0 mg
Glucose	100.0 mg

The above ingredients are dissolved in purified water to prepare 2 ml of a injection solution.

20

Pharmaceutical Preparation Example 6

Suppository comprising:

Compound of Example 1	100 mg
Witepsol <sup>®</sup> H 15	950 mg
Witepsol <sup>®</sup> E 75	950 mg

\* Witepsol is a trade name of Witten A-G in West Germany.

30

35 Pharmaceutical Preparation Example 7

Compound of Example 1	2 g
Ethyl para-hydroxybenzoate	0.025 g
Propyl para-hydroxybenzoate	0.015 g
Sodium laurylsulphonate	1.5 g
Propylene glycol	12.0 g
Stearyl alcohol	22.0 g
White vaseline	25.0 g

The above ingredients are dissolved in purified water to prepare 100.0 g of hydrophilic ointment.

Pharmacological Experiments:

50 Effect of the compounds on in vitro cell growth of various tumor cell lines:

The assay of cytotoxicity is dependent on the cellular reduction of MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] by the mitochondrial dehydrogenase of viable cells to a blue formazan product which can be measured spectrophotometrically. Cell suspension, 100  $\mu$ l, containing  $10^4$  tumor cells with or without test sample was incubated for 72 hr in 5% CO<sub>2</sub>-95% air at 37° C in 96-well microculture plates (Falcon 3072). The 10  $\mu$ l of MTT(5 mg/ml) in saline was added to each culture well and incubated for 4 hr at 37° C. The formazan crystals were solubilized by the addition of 100  $\mu$ l of 10% SDS/0.01 N HCl to each well, and incubated for 24 hr at 37° C. The plates were then read at 570 nm on the plate reader. The percentage of cytotoxicity was calculated from the formula:

$$\text{Cytotoxicity}(\%) = \left[ 1 - \frac{(\text{OD}_{570\text{nm}} \text{ of test well})}{(\text{OD}_{570 \text{ nm of control well}})} \right] \times 100$$

2'-Deoxy-2'-methylidene-5-fluorocytidine was dissolved in distilled water, and diluted in RPM1 1640 medium. Tumor cell lines were maintained in RPM1 1640 medium (Nissui Seiyaku Co., Ltd., Tokyo, Japan).


Table 1 below shows effect of 2'-deoxy-2'-methylidene-5-fluorocytidine on the growth of various tumor cell lines. IC<sub>50</sub> values gave the concentrations showing the 50% cytotoxicity.

Table 1

Effect of 2'-deoxy-2'-methylidene-5-fluorocytidine on the growth of various tumor cell lines	
Tumor cell line and its origin	IC <sub>50</sub> (μg/ml)
L1210/c; mouse leukemia	0.33
B16; mouse melanoma	8.9
CCRF-CEM; human T cell leukemia	0.14
MOLT4; human T cell leukemia	0.036
U937; human histiocytic leukemia	0.44
HL60; human promyelocytic leukemia	0.040
PC13; human lung large cell carcinoma	1.0

### Claims

(1) A 2'-methylidenepyrimidine nucleoside compound of the general formula




wherein R<sup>1</sup> stands for amino or hydroxy group; R<sup>2</sup> stands for a halogen or a lower alkyl when R<sup>1</sup> is amino or R<sup>2</sup> stands for an alkyl having 2 to 4 carbon atoms, an alkynyl having 2 to 4 carbon atoms or a haloalkyl when R<sup>1</sup> is hydroxy group; and R<sup>3</sup> stands for hydrogen or a phosphoric acid residue, or a salt thereof.

(2) A compound as claimed in claim (1) representable by the general formula



EP 0 360 018 A1




wherein R stands for a halogen, an alkyl having 2 to 4 carbon atoms, an alkynyl having 2 to 4 carbon atoms or haloalkyl and R<sup>3</sup> stands for hydrogen or a phosphoric acid residue, which is a tautomer of the compound as claimed in claim (1) wherein R<sup>1</sup> is hydroxy group.

20 (3) A compound as claimed in claim (1) which is 2'-deoxy-2'-methylidene-5-fluorocytidine or a salt thereof.

(4) An anticancer composition which is characterized by containing, as the active ingredient, a compound as claimed in claim (1).

25 (5) An anticancer composition as claimed in claim (4) wherein the active ingredient is 2'-deoxy-2'-methylidene-5-fluorocytidine or a salt thereof.

(6) A method for producing a 2'-methylidenepyrimidine nucleoside compound of the general formula :




wherein R<sup>1</sup> stands for amino or hydroxy group; R<sup>2</sup> stands for a halogen or a lower alkyl when R<sup>1</sup> is amino or R<sup>2</sup> stands for an alkyl having 2 to 4 carbon atoms, an alkynyl having 2 to 4 carbon atoms or a haloalkyl when R<sup>1</sup> is hydroxy group; and R<sup>3</sup> stands for hydrogen or a phosphoric acid residue, or a salt thereof which comprises ① in advance protecting the hydroxy groups at the 3'-and 5'-positions of the sugar moiety of a compound of the general formula :

55


469

90092670

EP 0 360 018 A1



wherein R' represents an alkyl having 2 to 4 carbon atoms, an alkynyl having 2 to 4 carbon atoms or a haloalkyl, oxidizing the hydroxy group at the 2'-position of the sugar moiety of said protected compound and introducing methylidene at the 2'-position, followed by deprotection of the protective groups of the hydroxy groups at 3'- and 5'-positions of the sugar moiety and, if desired, phosphorylation at the 5'-position of the sugar moiety or (2) in advance protecting the hydroxy groups at the 3'- and 5'-positions of the sugar moiety of a compound of the general formula :



wherein R'' represents a halogen or a lower alkyl, oxidizing the hydroxy group at the 2'-position of the sugar moiety of said protected compound and introducing methylidene at the 2'-position, followed by aminating 4-position in the base moiety of the thus-obtained methylidene compound to obtain the 4-amino compound, followed by deprotecting with 1,2,4-triazole to obtain the 4-(1,2,4-thiazol-1-yl) compound and converting the obtained compound to the 4-amino compound by passing ammonia gas thereto, followed by deprotecting reaction of the protective groups for the hydroxy groups and, if desired, phosphorylation.

(7) A method as claimed in Claim 6 wherein the reaction of methylidene compound with 1,2,4-triazol is conducted in the presence of phosphorus oxychloride and triethylamine.

(8) A method as claimed in Claim 6 wherein a 2'-methylidenepyrimidine nucleoside compound or a salt thereof is 2'-deoxy-2'-methylidene-5-fluorocytidine or a salt thereof.

90092670

470



European Patent  
Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 11 5516

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
A	CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 34, no. 4, April 1986, pages 1518-1523, Tokyo, JP; H. USUI et al.: "Synthesis of 8,2'-methano- and 8,2'-ethanoadenosines" * Page 1519, compound 3 *	1	C 07 H 19/06 A 61 K 31/70
A	CHEMICAL ABSTRACTS, vol. 102, no. 21, 27th May 1985, page 615, abstract no. 185424g, Columbus, Ohio, US; T. UEDA et al.: "Nucleosides and nucleotides. 51. Synthesis of 2'-C-nitromethyl derivatives of uridine and the structure of a carbon-bridged cyclonucleoside derived therefrom", & NUCLEOSIDES NUCLEOTIDES 1984, 3(2), 173-82 * Abstract *	1	
A	CHEMICAL ABSTRACTS, vol. 104, no. 17, 28th April 1986, page 742, abstract no. 149328m, Columbus, Ohio, US; T. UEDA et al.: "Synthesis of 2'-deoxy-6,2'-ethano-cyclouridine. (Nucleosides and nucleotides. Part 57), & NUCLEOSIDES NUCLEOTIDES 1985, 4(3), 401-9 * Abstract *	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 5)  C 07 H 19/00 A 61 K 31/00
P, X	CHEMICAL ABSTRACTS, vol. 110, no. 18, 1st May 1989, page 405, abstract no. 160409h, Columbus, Ohio, US; & JP-A-63 258 818 (YOSHITOMI PHARMACEUTICAL INDUSTRIES, LTD) 26-10-1988 * Abstract *	1-8	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-12-1989	Examiner SCOTT J.R.M.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>A : member of the same patent family, corresponding document</p>			



European Patent  
Office

# EUROPEAN SEARCH REPORT

Page 2

Application Number

EP 89 11 5516

## DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.9)
P, X	EP-A-0 310 673 (YAMASA SHOYO K.K.) * Abstract; page 1, line 36 - page 2, line 5; claims 1, 12 * -----	1-8	
			TECHNICAL FIELDS SEARCHED (Int. CL.5)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-12-1989	Examiner SCOTT J.R.M.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone  Y : particularly relevant if combined with another document of the same category  A : technological background  O : non-written disclosure  P : intermediate document</p> <p>T : theory or principle underlying the invention  E : earlier patent document, but published on, or after the filing date  D : document cited in the application  L : document cited for other reasons  A : member of the same patent family, corresponding document</p>			

EPO FORM 183 (1.1.87) (P.0401)

472

90092670

This Page is inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record

## BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images  
problems checked, please do not report the  
problems to the IFW Image Problem Mailbox**